

Individual assignment No 2

1. Calculate the final pressure when 1 mole of nitrogen at 300 K and 100 atm is heated at constant volume until attaining 500 K.
2. Calculate the pressure exerted by 1 mole of CO₂ behaving as (a) perfect gas and (b) van der Waals gas, when it is confined in the following conditions: T = 273.15 K and V = 22.414 L (constants of the van der Waals equation: a=3.592 atm·L²·mol⁻² and b=4.267·10⁻²·L·mol⁻¹)
3. Calculate the pressure for 1 mole of He at 25 °C to be in same volume with 1 mole of H₂ at 1 atm and 25 °C.
4. Calculate the pressure exerted by 1 mole of ethane (C₂H₆) behaving as: (a) perfect gas; (b) van der Waals gas, when confined in the following conditions: i) 273.15 K and V = 22.414 L; ii) 1000 K and 100 cm³. Data: a=5.489 L²·atm·mol⁻² and b=6.380·10⁻² L·mol⁻¹.
5. The mass percentage of dry air at sea level is approximately: N₂ – 75.5%; O₂ – 23.2%, and Ar – 1.3%. What are the partial pressures of each component when the total pressure is 1 atm?
6. A sample of 87 mg of an ideal gas at 0.600 bar duplicates its volume and triplicates its temperature. What will be the final pressure?
7. Two moles of an ideal gas at 500 K are isothermally and reversibly compressed until a final volume will equal to 1/10 of the initial volume. Calculate: (a) ΔU (b) ΔH (c) work done by the gas (d) heat absorbed by the gas. Specific molar isochoric heat capacity of gas should be taken according to MKT 2.5R.
8. One mole of an ideal gas expands from 10 L and 0 °C to 20 L and 100 °C. Taking C_V = 20 J·K⁻¹·mol⁻¹, calculate ΔU, W and Q for each of the following alternative steps for the whole process: (a) Isothermal and reversible expansion at 0 °C from 10 L to 20 L, followed by a constant volume heating until 100 °C and (b) Heating of 10 L at constant volume until 100 °C, followed by an isothermal and reversible expansion until 20 L.